

## Ecophysiological responses of *Eichhornia crassipes* (Mart.) Solms to As<sup>5+</sup> under different stress conditions

A.C. MENEGUELLI-SOUZA<sup>+</sup>, A.P. VITÓRIA, T.O. VIEIRA, M.S.O. DEGLI-ESPOSTI, and C.M.M. SOUZA

*Laboratório de Ciências Ambientais, Centro de Biociência e Biotecnologia, Universidade Estadual do Norte Fluminense, 28013 – 602, Campos dos Goytacazes, RJ, Brazil*

### Abstract

Arsenic is a critical contaminant that is released into the environment through geochemical processes and anthropic actions. Two independent hydroponic experiments were performed to evaluate the ecophysiological responses of water hyacinth [*Eichhornia crassipes* (Mart.) Solms] to As under various stress conditions. In experiment 1, water hyacinth was exposed to As<sup>5+</sup> at concentrations of 0, 0.2, 2.0, and 20 mg L<sup>-1</sup> for 0, 2, and 4 d; in experiment 2, water hyacinth was exposed at concentrations of 0, 0.025, 0.05, and 0.1 mg L<sup>-1</sup> for 0, 10, and 20 d. In both experiments, As accumulation in plant tissue was proportional to its increase in the nutrient solution; As concentrations were higher in roots than in shoots. Detrimental effects of As on gas exchange were observed and were more pronounced in experiment 1. In experiment 1, at the beginning on the second day of exposure, significant decreases of maximum photochemical efficiency of PSII ( $F_v/F_m$ ), variable chlorophyll fluorescence ( $F_v/F_0$ ), and photosynthetic pigment contents were observed in plants exposed to 2.0 and 20 mg(As<sup>5+</sup>) L<sup>-1</sup>. It indicated that damage to the photosynthetic apparatus had occurred. No changes in  $F_v/F_m$ ,  $F_v/F_0$ , and contents of photosynthetic pigments were observed in the plants grown in the presence of 0.2 mg(As<sup>5+</sup>) L<sup>-1</sup> (in experiment 1) or after any of the treatments in experiment 2, indicating plant tolerance. Elevated nonphotochemical quenching was observed in experiment 2 after 20 d of exposure to As; it was as a part of protection mechanisms of the photosynthetic apparatus in these plants. The results obtained here indicate that the use of water hyacinth for As<sup>5+</sup> removal from highly impacted environments is limited but that it is effective in remediating sites with a low contamination.

*Additional key words:* carotenoid; chlorosis; photosynthesis; remediation; senescence; trace element.

### Introduction

Arsenic is widespread in environment and is one of the most toxic contaminants to human health (Jain and Ali 2000). This metalloid has several oxidation states, but in natural waters, it is mostly present in inorganic form as As<sup>3+</sup> and As<sup>5+</sup>. Environmental problems related to As result from its mobilization under natural conditions (Schmöger *et al.* 2000) and from several anthropic activities, such as insecticide application and mining (Azcue and Nriagu 1995). When As comes from anthropic activities, its presence in the medium increases (Smedley and Kinniburgh 2001), generally resulting in toxic As

concentrations in aquatic and terrestrial ecosystems.

Some plant species are able to adapt to sites that are enriched in trace elements. Tolerance to trace elements can be constitutive or induced by exposure (Verkleij and Schat 1992). In general, plants are tolerant to As concentrations in the medium up to 50 mg L<sup>-1</sup> (Jiang and Sing 1994). However, plant species differ in their sensitivity to As. Arsenic phytotoxicity depends on several factors, such as its concentration in the medium, chemical species, absorption pathway, and bioavailability (Martínez-Sánchez *et al.* 2011). As can exhibit toxic effects on

Received 13 May 2015, accepted 2 July 2015, published as online-first 28 August 2015.

<sup>+</sup>Corresponding author; e-mail: [annalizameneguelli@hotmail.com](mailto:annalizameneguelli@hotmail.com)

*Abbreviations:* Car – carotenoids; Chl – chlorophyll; C<sub>i</sub> – intercellular CO<sub>2</sub> concentration; DMSO – dimethylsulfoxide; F<sub>0</sub> – minimal fluorescence yield of the dark-adapted state; F<sub>m</sub> – maximal fluorescence yield of the dark-adapted state; F<sub>v</sub> – variable fluorescence; F<sub>v</sub>/F<sub>0</sub> – variable chlorophyll fluorescence; F<sub>v</sub>/F<sub>m</sub> – maximum photochemical efficiency of PSII; g<sub>s</sub> – stomatal conductance; NPQ – nonphotochemical quenching; P<sub>N</sub> – net photosynthetic rate; q<sub>P</sub> – photochemical quenching.

*Acknowledgements:* The authors wish to thank the Coordination for the Improvement of Higher Education Personnel (CAPES), Research Foundation for the State of the Rio de Janeiro (FAPERJ), and the State University of the North Fluminense Darcy Ribeiro for financial support.

photosynthesis at the level of electron transport, CO<sub>2</sub> fixation, enzyme activities, photosynthetic pigments, and PSII functional activity (Päivöke and Simola 2001, Singh *et al.* 2009, Garg and Singla 2011). However, at low concentrations, tolerance mechanisms may be triggered, such as an increase in antioxidant enzyme activity, thermal dissipation (carotenoids and NPQ) and photochemical (q<sub>p</sub>) of excess energy, thereby avoiding the formation of singlet oxygen in the PSII reaction center (Burns *et al.* 2003, Vitória *et al.* 2010, 2011; Pereira *et al.* 2011).

However, some plant species, called hyperaccumulators, tolerate high concentrations of trace elements, such as As, in their tissues (Meharg and Hartley-Whitaker 2002). Ladder brake (*Pteris vittata*) was the first plant species reported to be the As hyperaccumulator, accumulating between 3,280 and 4,980 mg kg<sup>-1</sup> in its biomass (Ma *et al.* 2001). Water hyacinth (*Eichhornia crassipes*) is a free-floating macrophyte and is useful for the phytoremediation of contaminated aquatic environments due to its high biomass production and rapid propagation (Giri and Patel 2012).

As contents in the medium vary with the source of

## Materials and methods

**Plant material and As treatment:** Water hyacinths [*Eichhornia crassipes* (Mart.) Solms] of young and healthy appearance were collected at the Lagoa do Campelo, located in Campos dos Goytacazes, the state of Rio de Janeiro, Brazil (21°39'01"S, 41°11'W), during April 2011 (autumn) and January 2013 (summer). The plants were grown under greenhouse conditions in 10 L polypropylene pots containing 8 L of a nutrient solution (Hoagland and Arnon 1950, Smart and Barko 1985). The pots were distributed randomly. As was added in the form of Na<sub>2</sub>HAsO<sub>4</sub> (As<sup>5+</sup>). In experiment 1, the plants were subjected to 0, 0.20, 2.0, and 20 mg(As<sup>5+</sup>) L<sup>-1</sup> for 0, 2, and 4 d. Experiment 2 was designed according to results obtained in the experiment 1; the plants were subjected to 0, 0.025, 0.050, and 0.10 mg(As<sup>5+</sup>) L<sup>-1</sup> for 0, 10, and 20 d. All treatments were performed in triplicates (*n* = 3). Hydrogen potential, electrical conductivity, and dissolved O<sub>2</sub> were monitored throughout the experiments.

**As determination:** In order to determine As in plant tissues, the plants were harvested at the end of the experiments, separated into roots and shoots, and dried at 60°C for one week. When the plant material became dry, it was macerated in liquid nitrogen. Acid digestion was performed using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and a 1:1 mix of sulfuric and nitric acids (H<sub>2</sub>SO<sub>4</sub>:HNO<sub>3</sub>). The digestion was performed under pressure in a *Mars Xpress microwave* (CEM, USA). As was determined by inductively coupled plasma atomic emission spectroscopy (ICP-AES) using a *Liberty Series II* spectrometer (*Varian*, Australia). Values were expressed in mg(As) kg<sup>-1</sup> (dry mass, DM).

contamination and the environmental matrix to which the element is associated; it is higher in sediments, where it can reach between 161 and 4,709 mg(As) kg<sup>-1</sup> (Borba *et al.* 2003, Rezende *et al.* 2011). Lower As concentrations between 2 × 10<sup>-3</sup> and 3 mg L<sup>-1</sup> are found in the water column (Borba *et al.* 2004, Gonçalves *et al.* 2007). Permissible values in water, according to its use, were established to be between 1.4 × 10<sup>-4</sup> and 0.5 mg(As) L<sup>-1</sup> by the National Environmental Council (CONAMA 2005).

The goal of the present study was to determine and compare photosynthetic adaptation strategies of water hyacinth to As under a variety of contamination conditions and to evaluate the energy cost of tolerance mechanisms to the plant. A wide range of stress conditions was tested. With longer exposure times at concentrations close to the permissible levels established by the legislation, it is expected that the plant should be able to adapt its energy content to develop As tolerance mechanisms. In contrast, it is expected that short periods of exposure associated with high As concentrations do not allow the plants to maintain spending its resources on tolerance processes.

**Gas exchange:** Net photosynthetic rate (*P<sub>N</sub>*), stomatal conductance (*g<sub>s</sub>*), and substomatal CO<sub>2</sub> concentration (*C<sub>i</sub>*) were measured between 08:00 and 11:00 h using an infrared gas analyzer (*Ciras-2*, *PP Systems*, UK). Plants were measured under PPFD of 2,000 μmol m<sup>-2</sup> s<sup>-1</sup>, CO<sub>2</sub> concentration of 380 μmol mol<sup>-1</sup>, 25 ± 3°C, and 80% humidity. Measurements were performed in two intact and healthy leaves from each plant for all treatments (*n* = 3).

**Chlorophyll (Chl) *a* fluorescence** measurements were performed between 12:00 and 13:00 h using a modulated light portable fluorometer (*FMS2*, *Hansatech*, UK). Six leaves from each treatment (*n* = 3) were kept in the dark for 30 min and then exposed to a modulated beam of light [approximately 6 μmol(photon) m<sup>-2</sup> s<sup>-1</sup> at 660 nm], followed by a saturating pulse of white actinic light [10,000 μmol(photon) m<sup>-2</sup> s<sup>-1</sup>], which was applied for 0.8 s (adapted from Genty *et al.* 1989). Minimal fluorescence (*F<sub>0</sub>*), maximum fluorescence (*F<sub>m</sub>*), photochemical quenching (q<sub>p</sub>), and nonphotochemical quenching (NPQ) were measured. Variable fluorescence (*F<sub>v</sub>* = *F<sub>m</sub>* - *F<sub>0</sub>*), *F<sub>v</sub>*/*F<sub>m</sub>*, and *F<sub>v</sub>*/*F<sub>0</sub>* were calculated according to van Kooten and Snel (1990). The measurements were performed on the same leaves used for the gas-exchange measurements.

**Photosynthetic pigments:** Three leaf discs (*n* = 3) were collected from each treatment to quantify Chl *a* and *b* and carotenoids (Car). Discs were cut, placed in capped plastic tubes containing 5 mL of dimethyl sulfoxide (DMSO) as an organic solvent, and stored in the dark for five days. Samples were measured using a spectrophotometer (*Model*

TCC-240A, Shimadzu, Japan) at 665, 649, and 480 nm. Chl *a* and *b* and Car were calculated according to Wellburn (1994). Total Chl concentration and Chl *a/b* and total Chl/Car ratios were calculated. All laboratory work was performed under low light. All values are expressed in  $\mu\text{mol cm}^{-2}$ .

## Results

**Visual evaluation:** Signs of chlorosis and necrosis were observed in plants 4 d following the addition of 2.0 and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  (experiment 1) (Fig. 1C,D). In experiment 2, signs of chlorosis and necrosis were observed 20 d after addition of 0.025  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  (Fig. 1F).

**As concentration:** Higher concentrations were observed in roots compared to shoots in both experiments (Table 1). As




Fig. 1. Visual aspects of water hyacinth exposed to  $\text{As}^{5+}$ . *Left column* – experiment 1 (after 4 d of As exposure). *Right column* – experiment 2 (after 20 d of As exposure). *A* – control plant; *B, C, D* – plants exposed to 0.20, 2.0, and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$ , respectively. *E* – control plant; *F, G, H* – plants exposed to 0.025, 0.050, and 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$ , respectively.

**Statistical analysis:** As quantification, gas exchange, Chl *a* fluorescence, and photosynthetic pigment data were subjected to a two-way analysis of variance (ANOVA), and the means were compared using Tukey's test ( $p < 0.05$ ).

root to shoot translocation was higher in the plants subjected to 2.0  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  in experiment 1 (day 4) and for plants exposed to 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  in experiment 2 (day 20).

Table 1. Arsenic concentration [ $\text{mg kg}^{-1}(\text{DM})$ ] in water hyacinth exposed to 0, 0.20, 2.0, and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  for 4 d (experiment 1) and 0, 0.025, 0.050, and 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  for 20 d (experiment 2). Each value represents the mean  $\pm$  SD ( $n = 3$ ). *Uppercase letters*: comparing the same treatment in different parts of the plant. *Lowercase letters*: comparison of different treatments in the same part of the plant. Values followed by *different letters* differ significantly ( $p < 0.05$ ).

Treatment [ $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$ ]	As [ $\text{mg kg}^{-1}(\text{DM})$ ]	
	Shoots	Roots
Control	$0 \pm 0.00^{\text{Ac}}$	$2.66 \pm 3.31^{\text{Ac}}$
0.20	$51.14 \pm 29.62^{\text{Bb}}$	$524.83 \pm 176.24^{\text{Ab}}$
2.0	$1,626.80 \pm 540.00^{\text{Aa}}$	$968.31 \pm 178.09^{\text{Ab}}$
20	$1,841.66 \pm 501.67^{\text{Ba}}$	$2,915.28 \pm 55.39^{\text{Aa}}$
Control	$0 \pm 0.00^{\text{Ab}}$	$1.39 \pm 0.33^{\text{Ac}}$
0.025	$1.09 \pm 0.36^{\text{Bb}}$	$28.34 \pm 3.01^{\text{Ab}}$
0.050	$3.43 \pm 1.08^{\text{Bb}}$	$19.92 \pm 2.30^{\text{Ab}}$
0.10	$11.70 \pm 0.66^{\text{Ba}}$	$34.55 \pm 2.49^{\text{Aa}}$

**Gas-exchange parameters:** In experiment 1, the highest  $P_N$  decrease was observed after 2 d of exposure to 2.0 and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$ . This decrease was still observed on day 4 in these treatments (Fig. 2A). In addition, increased  $C_i$  and decreased  $g_s$  were observed for these treatments (Fig. 2B,C). The lowest  $P_N$  for experiment 2 was observed after 20 d of plant exposure to 0.05 and 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  (Fig. 2D), followed by an increase in  $C_i$  (Fig. 2E). However, the decrease in  $P_N$  was less pronounced for experiment 2 than that for experiment 1. A significant decrease in  $g_s$  was observed in experiment 2 after 20 d of As exposure in all treatments (Fig. 2F).

**Analysis of Chl *a* fluorescence parameters:** In experiment 1, significant decreases in  $F_v/F_m$  and  $F_v/F_0$  were observed in the presence of 2.0 and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  starting from 2 d after exposure (Table 2). On day 4,  $F_v/F_m$  varied between 0.86 (control) and 0.34 (20  $\text{mg L}^{-1}$ ), and  $F_v/F_0$  varied between 5.79 (control) and 0.51 (20  $\text{mg L}^{-1}$ ). Significant  $q_p$  decreases were observed in the presence of 2.0  $\text{mg L}^{-1}$  (0.84) and 20  $\text{mg L}^{-1}$  (0.86) starting after 2 d of exposure. In experiment 2,  $F_v/F_m$ ,  $F_v/F_0$ , and  $q_p$  were stable




Fig. 2. Effects of  $\text{As}^{5+}$  exposure on net photosynthetic rate (A,D), substomatal  $\text{CO}_2$  concentration (B,E), and stomatal conductance (C,F) in water hyacinth leaves. *Left column* – experiment 1. *Right column* – experiment 2. Plants were grown with 0 ( $\blacklozenge$ ), 0.20 ( $\square$ ), 2.0 ( $\blacktriangle$ ), and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  ( $\times$ ) for 0, 2, and 4 d and with 0 ( $\blacklozenge$ ) 0.025 ( $\blacktriangle$ ), 0.050 ( $\square$ ), and 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  ( $\times$ ) for 0, 10, and 20 d.  $P_N$  – net photosynthetic rate,  $C_i$  – intercellular  $\text{CO}_2$  concentration,  $g_s$  – stomatal conductance. Data points represent the averages  $\pm$  SD ( $n = 3$ ).

in all treatments and only varied significantly with time. In experiment 1, NPQ increased significantly in the presence of 2.0 and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  on day 2 and decreased significantly on day 4. In experiment 2, NPQ increased in the presence of 0.050 and 0.10  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  after 20 d following the addition. For both experiments,  $F_v/F_0$  was more sensitive than  $F_v/F_m$  in detecting variations due to the presence of  $\text{As}^{5+}$ .

## Discussion

The uptake and accumulation of As by several plant species are well documented (Ma *et al.* 2001, Robinson *et al.* 2003, Xue and Yan 2011, Farnese *et al.* 2014). In the present study, the accumulation of As in plant tissue and its translocation into the shoots were mostly determined by the As concentration in the growth medium, not by the duration of exposure. The higher retention of As in the roots, which was observed in the present study, might be due to (1) the ability of As to be bond to root cell wall

**Photosynthetic pigments:** In experiment 1, Chl *a* and *b* concentrations significantly decreased in the presence of 2.0 and 20  $\text{mg}(\text{As}^{5+}) \text{L}^{-1}$  (Table 3). The Chl *a/b* and total Chl/Car ratios decreased in the presence of 20  $\text{mg} \text{L}^{-1}$ . No changes in pigment concentrations were observed during experiment 2.

components (Salt *et al.* 1998), (2) a decrease in the conduction of As *via* water flow through the xylem (Stoeva *et al.* 2005), and (3) the complexation of As with thiol groups ( $\text{As}^{5+}$  is reduced to  $\text{As}^{3+}$  when it enters cells, Mathews *et al.* 2010). Arsenite has high affinity for thiol groups, which are responsible for its sequestration into vacuoles in roots (Zhao *et al.* 2009). The importance of roots as a preferential site of trace element accumulation has direct effects on photosynthesis, which is crucial for

Table 2. Chlorophyll *a* fluorescence parameters of the leaves in water hyacinth (dark-adapted) exposed to 0, 0.20, 2.0, and 20 mg(As<sup>5+</sup>) L<sup>-1</sup> for 0, 2, and 4 d and 0, 0.025, 0.050, and 0.10 mg(As<sup>5+</sup>) L<sup>-1</sup> for 0, 10, and 20 d. Each value represents the mean ± SD (*n* = 3). F<sub>v</sub>/F<sub>m</sub> – maximum quantum efficiency; F<sub>v</sub>/F<sub>0</sub> – variable chlorophyll fluorescence; qp – photochemical quenching; NPQ – nonphotochemical quenching. *Uppercase letters* indicate significant differences in the same treatment at different exposure times while *lowercase letters* differ treatments at the same time of exposure (*p*<0.05).

Time [d]	Treatment [mg(As <sup>5+</sup> ) L <sup>-1</sup> ]	F <sub>v</sub> /F <sub>m</sub>	F <sub>v</sub> /F <sub>0</sub>	qp	NPQ
0	Control	0.84 ± 0.01 <sup>Aa</sup>	5.43 ± 0.48 <sup>Aa</sup>	0.98 ± 0.01 <sup>Aa</sup>	0.08 ± 0.00 <sup>Aa</sup>
	0.20	0.86 ± 0.01 <sup>Aa</sup>	5.96 ± 0.41 <sup>Aa</sup>	0.94 ± 0.03 <sup>Aa</sup>	0.09 ± 0.00 <sup>Aa</sup>
	2.0	0.86 ± 0.01 <sup>Aa</sup>	6.21 ± 0.41 <sup>Aa</sup>	0.95 ± 0.04 <sup>Aa</sup>	0.09 ± 0.00 <sup>Ba</sup>
	20	0.85 ± 0.01 <sup>Aa</sup>	5.79 ± 0.41 <sup>Aa</sup>	0.94 ± 0.05 <sup>Aa</sup>	0.11 ± 0.00 <sup>Aa</sup>
2	Control	0.87 ± 0.01 <sup>Aa</sup>	6.55 ± 0.38 <sup>Aa</sup>	0.97 ± 0.01 <sup>Aa</sup>	0.06 ± 0.00 <sup>Aa</sup>
	0.20	0.86 ± 0.01 <sup>Aa</sup>	6.23 ± 0.35 <sup>Aa</sup>	0.97 ± 0.01 <sup>Aa</sup>	0.07 ± 0.00 <sup>Aa</sup>
	2.0	0.72 ± 0.09 <sup>Bb</sup>	2.94 ± 1.59 <sup>Bb</sup>	0.84 ± 0.05 <sup>Bb</sup>	0.29 ± 0.00 <sup>Ab</sup>
	20	0.68 ± 0.13 <sup>Bb</sup>	2.56 ± 1.24 <sup>Bb</sup>	0.86 ± 0.01 <sup>Bb</sup>	0.36 ± 0.00 <sup>Ab</sup>
4	Control	0.85 ± 0.01 <sup>Aa</sup>	5.79 ± 0.22 <sup>Aa</sup>	0.98 ± 0.02 <sup>Aa</sup>	0.13 ± 0.01 <sup>Aa</sup>
	0.20	0.84 ± 0.01 <sup>Aa</sup>	5.12 ± 0.53 <sup>Aa</sup>	0.98 ± 0.01 <sup>Aa</sup>	0.06 ± 0.00 <sup>Aa</sup>
	2.0	0.50 ± 0.05 <sup>Cb</sup>	0.99 ± 0.18 <sup>Bb</sup>	0.76 ± 0.08 <sup>Bb</sup>	0.01 ± 0.00 <sup>Ba</sup>
	20	0.34 ± 0.01 <sup>Cb</sup>	0.51 ± 0.03 <sup>Bb</sup>	0.86 ± 0.01 <sup>Bb</sup>	0.01 ± 0.00 <sup>Aa</sup>
0	Control	0.89 ± 0.01 <sup>Aa</sup>	8.28 ± 0.69 <sup>Aa</sup>	0.97 ± 0.02 <sup>Aa</sup>	0.09 ± 0.00 <sup>Aa</sup>
	0.025	0.89 ± 0.01 <sup>Aa</sup>	7.89 ± 0.84 <sup>Aa</sup>	0.95 ± 0.03 <sup>Aa</sup>	0.08 ± 0.00 <sup>Aa</sup>
	0.050	0.89 ± 0.01 <sup>Aa</sup>	8.02 ± 1.11 <sup>Aa</sup>	0.95 ± 0.02 <sup>Aa</sup>	0.06 ± 0.00 <sup>Ba</sup>
	0.10	0.89 ± 0.01 <sup>Aa</sup>	8.24 ± 0.53 <sup>Aa</sup>	0.94 ± 0.01 <sup>Aa</sup>	0.07 ± 0.00 <sup>Ba</sup>
10	Control	0.84 ± 0.01 <sup>Ba</sup>	5.56 ± 0.05 <sup>Ba</sup>	0.92 ± 0.03 <sup>Ba</sup>	0.06 ± 0.00 <sup>Aa</sup>
	0.025	0.85 ± 0.01 <sup>Ba</sup>	5.68 ± 0.30 <sup>Ba</sup>	0.92 ± 0.03 <sup>Aa</sup>	0.10 ± 0.00 <sup>Aa</sup>
	0.050	0.86 ± 0.00 <sup>Aa</sup>	6.21 ± 0.23 <sup>Ba</sup>	0.90 ± 0.02 <sup>Ba</sup>	0.08 ± 0.00 <sup>Ab</sup>
	0.10	0.86 ± 0.01 <sup>Aa</sup>	6.06 ± 0.37 <sup>Ba</sup>	0.89 ± 0.03 <sup>Ba</sup>	0.08 ± 0.00 <sup>Ba</sup>
20	Control	0.81 ± 0.02 <sup>Ba</sup>	5.45 ± 0.31 <sup>Ba</sup>	0.92 ± 0.01 <sup>Ba</sup>	0.09 ± 0.00 <sup>Aa</sup>
	0.025	0.81 ± 0.03 <sup>Ca</sup>	5.14 ± 0.15 <sup>Ba</sup>	0.91 ± 0.02 <sup>Aa</sup>	0.16 ± 0.00 <sup>Aa</sup>
	0.050	0.82 ± 0.04 <sup>Ba</sup>	4.83 ± 0.70 <sup>Ca</sup>	0.92 ± 0.02 <sup>ABa</sup>	0.17 ± 0.00 <sup>Aa</sup>
	0.10	0.81 ± 0.02 <sup>Ba</sup>	4.25 ± 0.42 <sup>Ca</sup>	0.92 ± 0.01 <sup>ABa</sup>	0.19 ± 0.00 <sup>Aa</sup>

Table 3. Chlorophyll (Chl) *a*, *b*, and carotenoids (Car) content, Chl *a/b* and total Chl/Car ratios in water hyacinth exposed to 0, 0.20, 2.0, and 20 mg(As<sup>5+</sup>) L<sup>-1</sup> for 4 d and 0, 0.025, 0.050, and 0.10 mg(As<sup>5+</sup>) L<sup>-1</sup> for 20 d. Each value represents the mean ± SD (*n* = 3). Values followed by *different letters* differ significantly (*p*<0.05).

Treatment [mg(As <sup>5+</sup> ) L <sup>-1</sup> ]	Chl <i>a</i>	Chl <i>b</i>	Total Chl	Car	Chl <i>a/b</i>	Total Chl/Car
Control	43.46 ± 4.24 <sup>A</sup>	14.70 ± 1.56 <sup>A</sup>	58.16 ± 5.75 <sup>A</sup>	8.88 ± 0.90 <sup>A</sup>	2.96 ± 0.09 <sup>A</sup>	6.55 ± 0.09 <sup>A</sup>
0.2	43.67 ± 2.07 <sup>A</sup>	14.33 ± 0.88 <sup>A</sup>	58.00 ± 1.61 <sup>A</sup>	9.79 ± 0.89 <sup>A</sup>	3.05 ± 0.30 <sup>A</sup>	5.97 ± 0.30 <sup>A</sup>
2.0	36.62 ± 0.00 <sup>B</sup>	10.95 ± 0.00 <sup>B</sup>	47.67 ± 0.00 <sup>B</sup>	8.10 ± 0.00 <sup>A</sup>	3.35 ± 0.35 <sup>A</sup>	5.87 ± 0.00 <sup>A</sup>
20	21.22 ± 5.79 <sup>B</sup>	8.42 ± 1.30 <sup>B</sup>	29.64 ± 7.02 <sup>C</sup>	8.19 ± 0.31 <sup>A</sup>	2.50 ± 0.34 <sup>B</sup>	3.13 ± 0.19 <sup>B</sup>
Control	50.70 ± 0.01 <sup>A</sup>	16.30 ± 0.00 <sup>A</sup>	66.96 ± 0.01 <sup>A</sup>	11.17 ± 0.00 <sup>A</sup>	3.11 ± 0.00 <sup>A</sup>	6.04 ± 0.01 <sup>A</sup>
0.025	54.85 ± 0.00 <sup>A</sup>	17.20 ± 0.00 <sup>A</sup>	72.05 ± 0.01 <sup>A</sup>	11.61 ± 0.00 <sup>A</sup>	3.20 ± 0.00 <sup>A</sup>	6.23 ± 0.00 <sup>A</sup>
0.05	46.63 ± 0.00 <sup>A</sup>	15.10 ± 0.00 <sup>A</sup>	62.72 ± 0.00 <sup>A</sup>	10.77 ± 0.00 <sup>A</sup>	3.10 ± 0.00 <sup>A</sup>	5.74 ± 0.00 <sup>A</sup>
0.10	49.05 ± 0.01 <sup>A</sup>	16.30 ± 0.00 <sup>A</sup>	65.40 ± 0.01 <sup>A</sup>	11.65 ± 0.00 <sup>A</sup>	3.00 ± 0.00 <sup>A</sup>	5.62 ± 0.00 <sup>A</sup>

biomass gain (Soltan and Rashed 2003). Visual symptoms of trace element phytotoxicity, such as necrosis (Singh *et al.* 2006) and chlorosis (Soltan and Rashed 2003), are related to the presence of these elements in shoot tissues and to decreases in essential nutrient uptake (Singh *et al.* 2005, Souza *et al.* 2009).

Significant changes in the physiological responses of vascular plants are caused by trace element accumulation

in their tissues (Jana 1988). The main mechanism responsible for the toxic effects of As is oxidative stress due to oxidation and reduction reactions that occur at the cellular level (Meharg and Hartley-Whitaker 2002). A marked drop in *P<sub>N</sub>* of maize plants was observed in the presence of 5 mg(Na<sub>3</sub>AsO<sub>4</sub>) L<sup>-1</sup> and was related to the decrease in *g<sub>s</sub>* (Stoeva *et al.* 2003). This finding was also observed in both experiments reported here. The direct effect of As on

$g_s$  might be due to the disruption of water uptake and transport that results from changes in roots during As uptake (Stoeva *et al.* 2003). Increased  $C_i$  and decreased  $g_s$  indicate that  $CO_2$  fixation is compromised both at the biochemical and ecophysiological levels (Stoeva and Bineva 2003, Gusman *et al.* 2013).

The decreased  $P_N$  due to the presence of  $As^{5+}$ , which was observed in experiment 1, might be caused by changes in PSII function (as indicated by the values of  $F_v/F_m$  and  $F_v/F_0$ ) and by the decrease in photosynthetic pigment concentrations. Values of  $F_v/F_m$  between 0.75 and 0.85 indicate that the photosynthetic apparatus is efficient (Bolhar-Nordenkamp *et al.* 1989), and the value of  $F_v/F_0$  lies in the range between 4 and 6 in healthy plants (Roháček 2002). Lower values than those above were observed in experiment 1 and in oat plants (*Avena sativa*) treated with high As concentrations (Stoeva and Bineva 2003). Decreases in the above ratios indicate photoinhibitory damage, which is caused by excess incident photon fluxes that occur in plants subjected to a wide variety of environmental stresses (Björkman *et al.* 1987).  $F_v/F_0$  has been suggested to be more sensitive in detecting small differences in PSII quantum yield compared to  $F_v/F_m$  (Oliveira *et al.* 2002, Paiva *et al.* 2009, Vitória *et al.* 2010), which was consistent with the results obtained in our experiment 1. Decreased values of  $F_v/F_m$  and  $F_v/F_0$  under  $As^{5+}$  stress might be due to decreased Chl concentration, as observed for *Jussiaea repens* in the presence of Cd (Li *et al.* 2008). The reactive oxygen species increase in the presence of As; these species might interact with Chl and oxidize it (Schützendübel and Polle 2002). The decrease in the Chl  $a/b$  ratio observed in experiment 1 indicated that Chl  $a$  was more affected than Chl  $b$ . A similar finding was observed for *Elodea densa* in the presence of Cu (Maleva *et al.* 2012). Changes in this ratio indicate modifications in thylakoid membranes that are reflected in the photosynthetic capacity (Päivöke and Simola 2001).

The fact that Chl  $a$  fluorescence and the photosynthetic pigment concentrations did not change in experiment 2 and in plants exposed to  $0.20\text{ mg}(As^{5+})L^{-1}$  (experiment 1) might be due to the low As concentration reported in the leaves (Table 1). Similar results were found in *Isatis cappadocia* and *Lactuca sativa* exposed to low As concentrations (Gusman *et al.* 2013, Karimi *et al.* 2013). Similarly, there was no significant change of the  $F_v/F_m$  ratio in the macrophytes, such as *Vallisneria gigantea*, *Azolla filiculoides*, and *Lemna minor*, which may be related to the low removal rate of As, ranging between 2 and 17% in this study (Iriel *et al.* 2015). The maintenance of the photosynthetic pigment concentrations explains the stability of  $F_v/F_m$  and  $F_v/F_0$  because changes in these parameters indicate decreased Chl concentrations or photooxidation (He *et al.* 2001). The high Chl  $a/b$  ratio indicated that the thylakoid stacking level was maintained (Table 3) (Lichtenthaler *et al.* 2006). The fact that this ratio and the total Chl/Car ratio did not change indicated that the pigments were resistant to the toxic elements (Maleva

*et al.* 2012).

The decrease of  $F_v/F_0$  may be the result not only of changes in the Chl concentration, as discussed above, but also due to  $F_0$  increase (not shown).  $F_0$  is the fluorescence intensity at the stage when all PSII reaction centers are open. When the Chl concentration is stable or reduced, the  $F_0$  increase may reflect the loss of efficiency of energy transfer from LHCs to PSII (Krause and Weis 1991).

To minimize the production of oxidant molecules in the photosynthetic apparatus, NPQ dissipates excess energy as heat, avoiding oxygen oxidation and protecting the plant against photooxidative damage (Li *et al.* 2000). In experiment 1, it was possible that the plants increased NPQ after 2 d of As exposure in an attempt to dissipate energy as heat and to protect the photosynthetic apparatus, as described largely under various types of stress (Lage-Pinto *et al.* 2008, Rabelo *et al.* 2013, Vieira *et al.* 2015). In some cases, the increase of heat dissipation (NPQ) can improve the plant performance and lead to a better adaptation to stress conditions. However, *E. crassipes* exposed for a short time (2 d) to Cr did not show this mechanism of energy dissipation (Paiva *et al.* 2009). Although *E. crassipes* increased NPQ when exposed to As for 2 d, this mechanism was unable to sustain the plant functionality for longer time (4 d, represented by the decrease of  $F_v/F_m$ ,  $F_v/F_0$ , and all gas-exchange parameters, Table 2, Fig. 2) and the senescence process was observed (Fig. 1). According to Demmig-Adams and Adams (1996), decreases in  $F_v/F_m$  under stress conditions are related to increases in heat dissipation (NPQ). Similarly, in experiment 2, the NPQ increase, which was observed after 20 d, reflected some of the protection mechanisms employed by the photosynthetic apparatus of these plants.

A positive correlation between NPQ and Car concentrations has been reported for water hyacinth in the field (Lage-Pinto *et al.* 2008), indicating that energy dissipation by the plants is a probable mechanism for the energy decrease observed in the electron transport chain. However, Car concentrations did not change in experiment 1. This finding might be related to the experimental conditions because there was no time for the plants to protect their photosynthetic apparatus from stress. Vitória *et al.* (2010) suggested that energy dissipation mechanisms are a long-term adaptation in water hyacinth plants. In experiment 2, it is possible that the increase in NPQ has been sufficient to dissipate the excess excitation energy at the PSII reaction center, thus rendering energy expenditure by the plant to increase the Car content unnecessary. Other tolerance mechanisms to oxidative stress caused by As might be activated, *e.g.*, an increase in the synthesis of thiols, such as phytochelatin, and increased antioxidant enzyme activity (Srivastava *et al.* 2007).

The results of hydroponic experiments are essential for understanding environmental contamination because these experiments allow the simulation of medium conditions in controlled assays. However, in hydroponics, the plant response varies with the conditions imposed, resulting in

punctual responses about particular stress. In the present study, experiments were performed in order to enable understanding of water hyacinth tolerance to various contamination conditions. Regarding the conditions tested in experiment 1, the exposure of water hyacinth to high As concentrations over short periods is more common when the source of contamination is anthropic, for example, due to toxic residues from metal industry. Regarding the conditions tested in experiment 2, water hyacinth exposure to low As concentrations over longer time periods are closer to As exposure originating from natural sources, such as the weathering of auriferous sulfide rocks. In

conclusion, our results indicated that the use of water hyacinth to remediate highly impacted environments is limited because significant damage to the photosynthetic apparatus was observed in experiment 1. In this experiment, reduction of stomatal conductance, decline in PSII function, and the concentration of photosynthetic pigments determined the stress caused by As. However, the As tolerance observed in experiment 2 and in plants exposed to 0.20 mg(As<sup>5+</sup>) L<sup>-1</sup> in experiment 1 indicated that As can be removed efficiently by water hyacinth in environments experiencing low contamination.

## References

- Azcue J.M., Nriagu J.O.: Impact of abandoned mine tailings on the arsenic concentrations in Moira Lake, Ontario. – *J. Geochem. Explor.* **52**: 81-89, 1995.
- Björkman O., Demmig B.: Photon yield of oxygen evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origin. – *Planta* **170**: 489-504, 1987.
- Bolhar-Nordenkamp H.R., Long S.P., Baker N.R.: Chlorophyll fluorescence as a probe of the photosynthetic competence of leaves in the field: a review of current instrumentation. – *Func. Ecol.* **3**: 497-514, 1989.
- Borba R.P., Figueiredo B.R., Rawlins B., Matschullat J.: Geochemical distribution of arsenic in waters, sediments and weathered gold mineralized rocks from Iron Quadrangle, Brazil. – *Environ. Geol.* **44**: 39-52, 2003.
- Borba R.P., Figueiredo B.R., Cavalcanti J.A.: [Arsenic in groundwater in Ouro Preto and Mariana, Iron Quadrangle (MG).] – *Geociências* **57**: 45-51, 2004. [In Portuguese]
- Burns J., Fraser P.D., Bramley P.M.: Identification and quantification of carotenoids, tocopherols and chlorophyll in commonly consumed fruits and vegetables. – *Phytochemistry* **62**: 939-947, 2003.
- CONAMA n° 357. March 17. Published in DOU n° 53, march 18. Section 1, 58-63, Ministério do Meio Ambiente, 2005. [In Portuguese]
- Demmig-Adams B., Adams W.W.: The role of xanthophyll cycle carotenoids in the protection of photosynthesis. – *Trend. Plant Sci.* **1**: 21-26, 1996.
- Farnese F.S., Oliveira J.A., Gusman G.S. *et al.*: Effects of adding nitroprusside on arsenic stressed response of *Pistia stratiotes* L. under hydroponic conditions. – *Int. J. Phyto-remediat.* **16**: 123-137, 2014.
- Garg N., Singla P.: Arsenic toxicity in crop plants: physiological effects and tolerance mechanisms. – *Environ. Chem. Lett.* **9**: 303-321, 2011.
- Genty B., Briantais J.M., Baker N.R.: The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. – *Biochim. Biophys. Acta* **990**: 87-92, 1989.
- Giri A.K., Patel R.K.: Phytoaccumulation potential and toxicity of arsenic ions by *Eichhornia crassipes* in hydroponic system. – *J. Bioremed. Biodegrad.* **3**: 2-6, 2012.
- Gonçalves J.A.C., Lena J.C., Paiva J.F. *et al.*: Arsenic in the groundwater of Ouro Preto (Brazil): its temporal behavior as influenced by the hydric regime and hydrogeology. – *Environ. Geol.* **53**: 785-793, 2007.
- Gusman G.S., Oliveira J.A., Farnese F.S. *et al.*: Arsenate and arsenite: the toxic effects on photosynthesis and growth of lettuce plants. – *Acta Physiol. Plant.* **35**: 1201-1209, 2013.
- He J., Lee S.K., Dodd I.C.: Limitations to photosynthesis of lettuce growth under tropical conditions: alleviation by root-zone cooling. – *J. Exp. Bot.* **52**: 1323-1330, 2001.
- Hoagland D.R., Arnon D.I.: *The Water Culture Method for Growing Plants without Soil*. Pp. 34. California Agriculture Station, Berkeley 1950.
- Iriel A., Dundas G., Cirelli A.F. *et al.*: Effect of arsenic on reflectance spectra and chlorophyll fluorescence of aquatic plants. – *Chemosphere* **119**: 697-703, 2015.
- Jain C.K., Ali I.: Arsenic: occurrence, toxicity and speciation techniques. – *Water Res.* **34**: 4304-4312, 2000.
- Jana S.: Accumulation of Hg and Cr by three aquatic species and subsequent changes in several physiological and bio-chemical plant parameters. – *Water Air Soil Poll.* **38**: 105-109, 1988.
- Jiang Q.Q., Singh B.R.: Effect of different forms and sources of arsenic on crop yield and arsenic concentration. – *Water Air Soil Poll.* **74**: 321-343, 1994.
- Karimi N., Shayesteh L.S., Ghasmpour H. *et al.*: Effects of arsenic on growth, photosynthetic activity, and accumulation in two new hyperaccumulating populations of *Isatis cappadocica* Desv. – *J. Plant Growth Regul.* **32**: 823-830, 2013.
- Krause G.H., Weis E.: Chlorophyll fluorescence and photosynthesis: the basics. – *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **42**: 313-349, 1991.
- Lage-Pinto F., Oliveira J.G., Cunha M.D. *et al.*: Chlorophyll a fluorescence and ultrastructural changes in chloroplast of water hyacinth as indicators of environmental stress. – *Environ. Exp. Bot.* **64**: 307-313, 2008.
- Lichtenthaler H.K., Kuhn G., Prenzel U. *et al.*: Chlorophyll-protein levels and degree of thylakoid stacking in radish chloroplasts from high-light, low-light and bentazon-treated plants. – *Physiol. Plantarum* **56**: 183-188, 2006.
- Li X.P., Björkman O., Shih C. *et al.*: A pigment-binding protein essential for regulation of photosynthetic light harvesting. – *Nature* **403**: 391-395, 2000.
- Li M., Zhang J., Ling T. *et al.*: Ecophysiological responses of *Jussiaea rapens* to cadmium exposure. – *Aquat. Bot.* **88**: 347-352, 2008.
- Ma L.Q., Komar K.M., Tu C. *et al.*: A fern that hyperaccumulates arsenic. – *Nature* **409**: 579-579, 2001.
- Maleva M.G., Nekrasova G.F., Borisova G.G. *et al.*: Effect of heavy metals on photosynthetic apparatus and antioxidant status of *Elodea*. – *Russ. J. Plant Physiol+* **59**: 190-197, 2012.
- Martínez-Sánchez M.J., Martínez-López S., García-Lorenzo



- M.L. *et al.*: Evaluation of arsenic in soils and plant uptake using various chemical extraction methods in soils affected by old mining activities. – *Geoderma* **160**: 535-541, 2011.
- Mathews S., Ma L.Q., Rathinasabapathi B. *et al.*: Arsenic transformation in the growth media and biomass of hyperaccumulator *Pteris vittata* L. – *Bioresource Technol.* **101**: 8024-8030, 2010.
- Meharg A.A., Hartley-Whitaker J.: Arsenic uptake and metabolism in arsenic resistant and nonresistant plant species. – *New Phytol.* **154**: 29-43, 2002.
- Oliveira J.G., Alves P.L.C.A., Magalhães A.C.N.: The effect of chilling on the photosynthetic activity in coffee (*Coffea arabica* L.) seedlings. The protective action of chloroplastid pigments. – *Braz. J. Plant Physiol.* **14**: 95-104, 2002.
- Paiva L.B., Oliveira J.G., Azevedo R.A. *et al.*: Ecophysiological responses of water hyacinth exposed to Cr<sup>3+</sup> and Cr<sup>6+</sup>. – *Environ. Exp. Bot.* **65**: 403-409, 2009.
- Päivöke A.E.A., Simola L.K.: Arsenate toxicity to *Pisum sativum*: mineral nutrients, chlorophyll content and phytase activity. – *Ecotox. Environ. Safe.* **49**: 111-121, 2001.
- Pereira F.J., Castro E.M., Oliveira C. *et al.*: [Anatomical and physiological mechanisms of water hyacinth plants to arsenic contamination tolerance.] – *Planta Daninha* **29**: 259-267, 2011. [In Portuguese]
- Rabelo G.R., Vitória A.P., da Silva M.V.A. *et al.*: Structural and ecophysiological adaptations to forest gaps. – *Trees-Struct. Funct.* **27**: 259-272, 2013.
- Rezende P.S., Moura P.A.S., Durão W.A.J. *et al.*: Arsenic and mercury mobility in Brazilian sediments from the São Francisco River Basin. – *J. Braz. Chem. Soc.* **22**: 910-918, 2011.
- Robinson B., Duwing C., Bolan N. *et al.*: Uptake of arsenic by New Zealand watercress (*Lepidium sativum*). – *Sci. Total Environ.* **301**: 67-73, 2003.
- Roháček K.: Chlorophyll fluorescence parameters: the definitions, photosynthetic meaning and mutual relationships. – *Photosynthetica* **40**: 13-29, 2002.
- Salt D.E., Smith R.D., Raskin I.: Phytoremediation. – *Annu. Rev. Plant Phys.* **49**: 643-668, 1998.
- Schmöger M., Oven M., Grill E.: Detoxification of arsenic by phytochelatins in plants. – *Plant Physiol.* **122**: 793-802, 2000.
- Schützendübel A., Polle A.: Plant responses to abiotic stress: heavy metal-induced oxidative stress and protection by mycorrhization. – *J. Exp. Bot.* **53**: 1351-1365, 2002.
- Singh N.K., Pandey G.C., Rai U.N. *et al.*: Metal accumulation and ecophysiological effects of distillery effluent on *Potamogeton pectinatus* L. – *Bull. Environ. Contam. Toxicol.* **74**: 857-863, 2005.
- Singh N., Ma L.Q., Srivastava M. *et al.*: Metabolic adaptations to arsenic-induced oxidative stress in *Pteris vittata* L. and *Pteris ensiformis* L. – *Plant Sci.* **170**: 274-282, 2006.
- Singh N., Ma L.Q., Va J.C. *et al.*: Effects of arsenic on nitrate metabolism in arsenic hyperaccumulating and non-hyperaccumulating ferns. – *Environ. Pollut.* **157**: 2300-2305, 2009.
- Smart R.M., Barko J.W.: Laboratory culture of submersed freshwater macrophytes on natural sediments. – *Aquat. Bot.* **21**: 251-263, 1985.
- Smedley P.L., Kinniburgh D.G.: Sources and Behavior of Arsenic in Natural Water. United Nations Synthesis Report on Arsenic in Drinking Water. Pp. 61. WHO, Geneva 2001.
- Soltan M.E., Rashed M.N.: Laboratory study on the survival of water hyacinth under several conditions of heavy metal concentrations. – *Adv. Environ. Res.* **7**: 321-334, 2003.
- Souza V.L., Silva D.C., Santana K.B. *et al.*: Effects of cadmium on the anatomy and photosynthesis of two aquatic macrophytes. – *Acta Bot. Bras.* **23**: 342-354, 2009.
- Srivastava S., Mishra S., Tripathi R.D. *et al.*: Phytochelatins and antioxidant systems respond differentially during arsenite and arsenate stress in *Hydrilla verticillata* (L.f.) Royle. – *Environ. Sci. Technol.* **41**: 2930-2936, 2007.
- Stoeva N., Bineva T.: Oxidative changes and photosynthesis in oat plants grown in As-contaminated soil. – *Bulgarian J. Agric. Sci.* **29**: 87-95, 2003.
- Stoeva N., Berova M., Zlatev Z.: Physiological response of maize to arsenic contamination. – *Biol. Plantarum* **47**: 449-452, 2003.
- Stoeva N., Berova M., Zlatev Z.: Effect of arsenic on some physiological parameters in bean plants. – *Biol. Plantarum* **49**: 293-296, 2005.
- van Kooten O., Snel J.F.H.: The use of chlorophyll fluorescence nomenclature in plant stress physiology. – *Photosynth. Res.* **25**: 147-150, 1990.
- Verkley J.A.C., Schat H.: Mechanisms of metal tolerance in higher plants. – In: Shaw A.J. (ed.): *Heavy Metal Tolerance in Plants: Evolutionary Aspects*. Pp. 179-193. CRC Press, New York 1992.
- Vieira T.O., Degli-Espositi M.S.O., Souza G.M. *et al.*: Photoacclimation capacity in seedling and sapling of *Siparuna guianensis* (Siparunaceae): Response to irradiance gradient in tropical forest. – *Photosynthetica* **53**: 11-22, 2015.
- Vitória A.P., Lage-Pinto F., Campaneli L.B. *et al.*: Ecophysiological adaptation and metal accumulation in water hyacinth from two tropical rivers. – *Braz. J. Plant. Physiol.* **22**: 49-59, 2010.
- Vitória A.P., Lage-Pinto F., Campaneli L.B. *et al.*: Structural and ecophysiological alterations of the water hyacinth [*Eichhornia crassipes* (Mart.) Solms] due to anthropogenic stress in Brazilian rivers. – *Braz. Arch. Biol. Technol.* **54**: 1059-1068, 2011.
- Vitória, A.P., Santos, J.L.S., Salomão, M.S.M. B. *et al.*: Influence of ecologic type, seasonality, and origin of macrophyte in metal accumulation, anatomy and ecophysiology of *Eichhornia crassipes* and *Eichhornia azurea*. – *Aquat. Bot.* **125**: 9-16, 2015.
- Xue P.Y., Yan C.Z.: Arsenic accumulation and translocation in the submerged macrophyte *Hydrilla verticillata* (L.f.) Royle. – *Chemosphere* **85**: 1176-1181, 2011.
- Wellburn A.R.: The spectral determination of chlorophylls *a* and *b*, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. – *J. Plant Physiol.* **144**: 307-313, 1994.
- Zhao F.J., Ma J.F., Meharq A.A. *et al.*: Arsenic uptake and metabolism in plants. – *New Phytol.* **181**: 777-794, 2009.